

GENERALISED MASS VARIATION FORMULA FOR A STATIONARY AXISYMMETRIC STAR OR BLACK HOLE WITH SURROUNDING ACCRETION DISC

Brandon Carter

G.A.R., Observatoire de Paris, 92 Meudon.

Abstract *The general relativistic mass-energy variation formula for axisymmetric equilibrium states of a selfgravitating system is developed in the particular case for which the relevant matter consists of a perfectly conducting multiconstituent fluid (or superfluid) with stationary circular or convective motion.*

1. Introduction

This communication presents an extension of the study¹ of mass variation formulae for stationary axisymmetric systems – relativistic stars and black holes with surrounding material discs – that is contained in my contribution (chapter 6) to the Einstein Centenary Survey: General Relativity, ed S.W. Hawking and W. Israel (Cambridge U.P. 1979); all numbered references here refer to equations presented therein.

The present work starts from the general formula

$$\delta M = \frac{1}{8\pi} \delta \int G_a{}^b k^a d\Sigma_b - \frac{1}{8\pi} \int G^{cd} h_{cd} k^a d\Sigma_a + \Omega^H \delta J_H + \frac{\kappa}{8\pi} \delta \mathcal{A} \quad (1)$$

(equation (6.342)¹) for the variation of the total asymptotically defined mass M of the system between two neighbouring stationary axisymmetric equilibrium states, where Ω^H , J_H , κ , $\mathcal{A}$ are the angular velocity, angular momentum, decay parameter (“surface gravity”) and surface area of the central black hole – if it is present – while G_{ab} is the Einstein tensor, h_{cd} the variation of the metric tensor, k^a the stationary Killing vector, and the integration is taken over a spacelike hypersurface Σ extending out to the asymptotically flat region with inner boundary on the black hole if it is present.

2. Electromagnetic and material contributions

We now introduce the energy momentum tensor T_{ab} and the corresponding momentum vector $\Pi^a = -T^a{}_b k^b$ (decomposed into electromagnetic and purely material parts, labelled by suffices _F and _M) as follows. When we substitute from the Einstein equations into the fundamental kinematic mass variation formula (1) we obtain

$$\delta M - \Omega^H \delta J_H - \frac{\kappa}{8\pi} \delta \mathcal{A} = \delta \int T^a{}_b k^b d\Sigma_a - \frac{1}{2} \int T^b{}_c h^c{}_b k^a d\Sigma_a \quad (2)$$

i.e.

$$\begin{aligned} \delta M - \Omega^H \delta J_H - \frac{\kappa}{8\pi} \delta \mathcal{A} + \delta \int \Pi_F^a \{k\} d\Sigma_a + \frac{1}{2} \int T_F^b{}_c h^c{}_b k^a d\Sigma_a \\ = -\delta \int \Pi_M^a \{k\} d\Sigma_a - \frac{1}{2} \int T_M^b{}_c h^c{}_b k^a d\Sigma_a \end{aligned} \quad (3)$$

To evaluate the electromagnetic contribution we first use the analogue of (6.306) for the momentum flux of a stationary field, namely

$$-\Pi_F^a = \Lambda_F k^a + (k^b A_b) j^a + \nabla_b (A_c k^c D_F^{ba}) \quad (4)$$

and then work out

$$\begin{aligned} |g|^{-1/2} \delta(|g|^{1/2} \Lambda_F k^a) &= \frac{1}{2} (T_F^b{}_c h^c{}_b - D_F^{bc} \delta F_{bc}) k^a \\ &= \left(\frac{1}{2} T_F^{bc} h_{bc} - j^b \delta A_b \right) k^a + 2 \nabla_b (D_F^{c[b} \delta A_c k^{a]}) , \end{aligned} \quad (5)$$

where the electromagnetic action density and displacement tensor are given – by (6.193) and (6.194)¹ – as $\Lambda_F = (1/16\pi) F_{ab} F^{ba}$ and $D_F^{ab} = (1/4\pi) F^{ab}$.

With the aid of the Green theorem (and using the asymptotic flatness conditions to eliminate a surface contribution at infinity) we are thus able to obtain

$$\begin{aligned} \delta \int_\Sigma \Pi_F^a d\Sigma_a - \frac{1}{2} \int_\Sigma T_F^b{}_c h^c{}_b k^a d\Sigma_a &= \int_\Sigma k^b A_b \delta(j^a d\Sigma_a) + \frac{1}{2} \int_\Sigma k^{[b} j^{a]} (\delta A_b) d\Sigma_a \\ &\quad - \frac{1}{2} \oint_S \frac{1}{2} k^c A_c D_F^{ba} dS_{ab} - \oint_S D_F^{c[b} (\delta A_c) k^{a]} dS_{ab} \end{aligned} \quad (6)$$

3. Perfectly conducting fluid

The result so far does not depend on the particular field equations satisfied by any matter that is present. In order to show the relationship with the first law of ordinary thermodynamics we shall now consider the explicit form of the right hand side of (3) in a particularly ideal case namely that of a *perfectly conducting multiconstituent fluid medium*. Previous descriptions of the first law by Bardeen 1973², Bardeen, Carter and Hawking 1973³ have been restricted to the more specialised case of a *non-conducting* medium, electromagnetic effects being taken into account in the more extended treatment given by Carter 1973b⁴.

The more general treatment presented here is based on the relativistic theory of a multiconstituent perfect fluid obtained as the natural extension of the two constituent case discussed by Carter 1976⁵. According to this theory (which includes the Landau-London two-fluid model for superfluidity as a special case in the non-relativistic limit) the material equations of motion are derivable from a Lagrangian scalar Λ_M which is a covariant function of a set of conserved current vectors $\vec{n}_X$ representing the fluxes of a set of independent constituents labelled by an index X , one of the constituents being usually (except in zero-temperature models) the *entropy*. (Thus in a typical application X might take the values I, II, III with $\vec{n}_I$ and $\vec{n}_{II}$ representing fluxes of positively charged

ions and negatively charged electrons respectively, while $\vec{n}_{\text{III}}$ is identified with an entropy flux vector $\vec{s}$.

Under the influence of arbitrary infinitesimal variations of the fluxes $\vec{n}$ and of the metric $\underline{g}$ the function Λ_{M} will have an infinitesimal variation of the form

$$\delta\Lambda_{\text{M}} = \mu_a^x \delta n_x^a + \frac{1}{2} \mu_a^x n_x^b h_b^a. \quad (7)$$

The covectors $\underline{\mu}^x$ so defined may appropriately be described as the *chemical potential one-forms* of the corresponding conjugate constituents. (In particular the covector conjugate to the entropy flux vector $\vec{s}$ may appropriately be denoted by $\underline{\Theta}$ and interpreted as a *temperature form*.) Their constituent indices are written upstairs in view of the fact that they transform *contravariantly* with respect to the fluxes under a change of *chemical basis* in the vector space of constituents.

When the medium interacts with an electromagnetic field the *total* Lagrangian will be taken to have the simple additive form

$$\Lambda = \Lambda_{\text{M}} + A_a j^a + \Lambda_{\text{F}} \quad (8)$$

where Λ_{F} is the electromagnetic Lagrangian (given by (6.193)¹) and where the electromagnetic current j^a appearing in the interaction term is given by

$$\vec{j} = e^x \vec{n}_x \quad (9)$$

where e^x represents the mean *electric charge per particle* of the x th constituent. Like the chemical potentials, the charges behave as components of a contravariant vector in the chemical constituent space. (It would in fact be easy to work with a more general Lagrangian involving a more intimate relationship between the fluxes $\vec{n}_x$ and the field $\underline{F}$. The restriction to the simple additive form (8) – which means that polarisability effects are excluded – has been made merely to facilitate the interpretation of the formalism.) Variation of the total Lagrangian now gives

$$\begin{aligned} \delta\Lambda \equiv & \pi_a^x \delta n_x^a + (j^a - j_{\text{F}}^a) \delta A_a - \nabla_c (D_{\text{F}}^{cb} \delta A_b) \\ & + \frac{1}{2} \left(n_x^a \pi_b^x - D_{\text{F}}^{ca} \nabla_b A_c + \nabla_c (D_{\text{F}}^{ca} A_b) + (j_{\text{F}}^a - j^a) A_b \right) h_a^b \end{aligned} \quad (10)$$

where we have used the abbreviation

$$j_{\text{F}}^a \equiv \nabla_b D_{\text{F}}^{ab} \quad (11)$$

in accordance with the scheme introduced in section 6.4¹, and where we introduce the notation

$$\underline{\pi}^x = \underline{\mu}^x + e^x \underline{A} \quad (12)$$

for what will play the role of an effective *four-momentum per particle* for the corresponding constituent. If we required the volume integral of Λ to be invariant under *arbitrary* variations of the fields $\underline{A}$ and $\vec{n}_x$ we would obtain not only the correct equations of motion

$$\vec{j}_{\text{F}} = \vec{j} \quad (13)$$

for the electromagnetic field, but also the too restrictive requirement that the momenta $\underline{\pi}^x$ vanish altogether. To obtain reasonable equations of motion for the chemical constituents one must demand invariance not under arbitrary current variations but only under variations of the restricted class that arise from *infinitesimal displacements of the world lines* of the idealised particles, or directly from variations of g_{ab} . Now it can easily be seen (e.g using the formulae given by Carter 1973a⁶) that an infinitesimal displacement $\vec{\xi}_x$ of the world lines of a conserved flux of particles (acting in conjunction with a variation $\underline{h}$ of the metric) induces a corresponding variation

$$\delta n^a = -\frac{1}{2}n^a h^c_c - n^a \nabla_b \xi^b + n^b \nabla_b \xi^a - \xi^b \nabla_b n^a \quad (14)$$

in the corresponding current vector $\vec{n}$.

The equations (7) to (13) were *chemically covariant* (i.e. unaffected by changes in the chemical basis of the space of constituents) in *form* although not necessarily in content (since the *specific* representation of the Lagrangian will in general depend on the chemical basis). However the choice of a particular set of chemical constituents implies the choice of a *preferred basis* in the vector space of constituents, so that the system of equations that results will not have even the appearance of chemical covariance. It is also to be remarked that although the identities derived in section 6.4.1¹ (which depended only on the covariance of the Lagrangian under general co-ordinate transformations of the space-time manifold as opposed to the chemical constituent vector space) will still be formally valid for the material lagrangian Λ_M – indeed (6.135)¹ has been implicitly used in the derivation of the form (7) – and for the total Lagrangian Λ , the quantities involved cannot be given the same direct interpretation (as representing energy-momentum etc. ...) for a *restrained* variational theory as in the free field theory for which they were originally intended. In the present theory the appropriate energy-momentum tensor cannot be obtained directly from the analogue of (6.138)¹ but can be read out from the expression

$$|g|^{-1/2} \delta(|g|^{1/2} \Lambda) \equiv T^{ab} h_{ab} + (j^a - j_F^a) \delta A_a - f_a^x \xi_x^a + \nabla_a \left(2\pi_b^x n_x^{[a} \xi_x^{b]} - A_b D_F^{ab} \right) \quad (15)$$

where the $\vec{\xi}_x$ (which do *not* behave as components of a well behaved vector under chemical basis transformations) are a set of arbitrary displacement vector fields acting on the world lines of the corresponding fluxes, and where the fields $\underline{f}^x$ (to which the same remark applies) are to be interpreted as generalised *4-force densities* acting on the particles of the corresponding constituents. One finds that the force covectors have the explicit form

$$\underline{f}^x \equiv \vec{n}_{|x|} \cdot \partial \underline{\pi}^x + \underline{\pi}^x \nabla \cdot \vec{n}_{|x|} \quad (16)$$

where the inclusion of the chemical indices within *straight bars* is introduced to denote suspension of the summation convention.

Each of the variational field equations

$$\underline{f}^x = 0 \quad (16)$$

can be decomposed by first contracting with the corresponding flux $\vec{n}^x$, to give *first* the conservation law

$$\nabla \cdot \vec{n}_x = 0 \quad 18$$

for each constituent, and *then* the canonical equation of motion

$$\vec{n}_{|x|} \cdot \partial \underline{\pi}^x = 0 \quad (19)$$

which determines the acceleration of the corresponding world lines. This last equation may be decomposed into a material and an electromagnetic part in the form

$$\vec{n}_{|x|} \cdot \partial \underline{\mu}^x = \underline{F} \cdot \vec{j}_x \quad (20)$$

where

$$\vec{j}^x = e^x \vec{n}_{|x|} \quad (21)$$

is the electric current density associated with the x th constituent. It is to be remarked that the condition (19) is precisely what is required for the flux of generalised vorticity

$$\underline{w}^x = \partial \underline{\pi}^x \quad (22)$$

to be conserved along the flow lines in the strong sense discussed in section 6.4.5¹, meaning that

$$(\alpha \vec{n}_{|x|}) \mathcal{L} \underline{w}^x = 0 \quad (23)$$

for an *arbitrary* scalar field α , which implies that the *flux* $\underline{w}^x$ over any 2-surface, which by Stokes theorem is equal to the *circulation*

$$\mathcal{C}^x = \oint_c \underline{\pi}^x \cdot d\vec{x} \quad (24)$$

around its boundary circuit c say, is conserved in the sense of being unchanged when the circuit (or the original 2-surface enclosed by it) is dragged at an arbitrary rate along the flow lines.

Unlike the field equations themselves, the expression for the *energy momentum* that is read out from (15) has a chemically covariant form. After subtracting out the Maxwellian contribution in accordance with the decomposition (6.299)¹ we are left with a *material* contribution to the energy-momentum given by

$$T_{\text{M}}^a{}_b = n_x^a \mu_b^x - \Psi_{\text{M}} g_b^a \quad (25)$$

where Ψ_{M} is a *thermodynamic potential density* that is related to the material Lagrangian by

$$\Psi_{\text{M}} = n_x^a \mu_a^x - \Lambda_{\text{M}} . \quad (26)$$

Provided that the electromagnetic field equations (13) are satisfied, the expression (6.299)¹ of the total metric energy momentum tensor as the sum of metric contributions T_{F}^{ab} and T_{M}^{ab} can be replaced by an expression in terms of canonical contributions in the form

$$T_b^a = \mathcal{T}_{\text{MC}}^a{}_b + \mathcal{T}_{\text{F}}^a{}_b + \nabla_c (D_{\text{F}}^{ca} A_b) \quad (27)$$

where $\mathcal{T}_{\text{F}}^a{}_b$ are components of the canonical electromagnetic energy momentum tensor introduced in (6.145)¹, and where the corresponding *canonical* material energy momentum tensor is defined by

$$\mathcal{T}_{\text{MC}}^a{}_b = n_x^a \pi_b^x - \Psi_{\text{M}} g_b^a . \quad (28)$$

Subject to the field equations, the *total* energy momentum tensor will of course be preserved, i.e.

$$\nabla_b T^b_a = 0 \quad (29)$$

and hence the separate parts will satisfy

$$\nabla_b T^b_{M a} = F_{ab} j^b = -\nabla_b T^b_{F a} , \quad (30)$$

$$\nabla_b \mathcal{T}^b_{MC a} = j^b \nabla_a A_b = -\nabla_b \mathcal{T}^b_{F a} . \quad (31)$$

For any vector field $\vec{\xi}$ we define associated canonical and metric energy - momentum flux vectors analogous to those introduced in section 6.4¹ by

$$\vec{\mathcal{P}}\{\xi\} = \vec{\mathcal{P}}_{MC}\{\xi\} + \vec{\mathcal{P}}_F\{\xi\} , \quad \mathcal{P}^a_{MC}\{\xi\} = -\mathcal{T}^a_{MC b} \xi^b . \quad (32)$$

$$\vec{\Pi}\{\xi\} = \vec{\Pi}_M\{\xi\} + \vec{\Pi}_F\{\xi\} , \quad \Pi^a_M\{\xi\} = -T^a_{M b} \xi^b , \quad (33)$$

the total being related by

$$\mathcal{P}^a\{\xi\} = \Pi^a\{\xi\} + \nabla_b (D^{ba}_F A_c \xi^c) . \quad (34)$$

It is obvious from (29) that the *total* metric energy - momentum fluxes will be conserved whenever $\vec{\xi}$ is a Killing vector, and hence that the same applies to the canonical energy - momentum flux, i.e.

$$\vec{\xi} \mathcal{L} \underline{g} = 0 \quad \Rightarrow \quad \begin{cases} \nabla \cdot \vec{\Pi}\{\xi\} = 0 \\ \nabla \cdot \vec{\mathcal{P}}\{\xi\} = 0 . \end{cases} \quad (35)$$

$$(36)$$

However the *canonical* energy - momentum flux also has the property that its *material part* will be conserved *separately* whenever the fields are *themselves invariant*. Clearly the equation of motion (19) implies (by the Cartan formula (6.20)¹) that we shall have a generalised Bernoulli theorem to the effect that

$$\vec{\xi} \mathcal{L} \underline{\pi}^x = 0 \quad \Rightarrow \quad \vec{n}_{|X|} \cdot \nabla (\vec{\xi} \cdot \underline{\pi}^x) = 0 , \quad (37)$$

i.e. that $\vec{\xi} \cdot \underline{\pi}^x$ is a *constant of the motion*, and hence, using (18) we see that

$$\left. \begin{aligned} \vec{\xi} \mathcal{L} \underline{\pi}^x = 0 , \quad \vec{\xi} \mathcal{L} \Psi_M = 0 \\ \vec{\xi} \mathcal{L} \underline{g} = 0 \end{aligned} \right\} \quad \Rightarrow \quad \nabla \cdot \vec{\mathcal{P}}_{MC}\{\xi\} = 0 . \quad (38)$$

This is to be compared with the corresponding result (6.157)¹ for the electromagnetic component, which tells us that the same result would follow from purely electromagnetic invariance, i.e.

$$\left. \begin{aligned} \vec{\xi} \mathcal{L} \underline{A} = 0 \\ \vec{\xi} \mathcal{L} \underline{g} = 0 \end{aligned} \right\} \quad \Rightarrow \quad \nabla \cdot \vec{\mathcal{P}}_{MC}\{\xi\} = 0 . \quad (39)$$

It may be noticed that across any *invariant* hypersurface, i.e. one whose surface element satisfies

$$\vec{\xi} \cdot d\underline{\Sigma} = 0 \quad (40)$$

the canonical material energy - momentum flux is interpretable as the *total* rate of transport of the Bernoulli quantity, i.e.

$$\mathcal{P}_{\text{MC}}^a \{\xi\} d\Sigma_a = (\vec{\xi} \cdot \underline{\pi}^x) dN_x \quad (41)$$

where dN_x denotes the number flux of the x th constituent across $d\underline{\Sigma}$, i.e.

$$dN_x = -\vec{n}_x \cdot d\underline{\Sigma} . \quad (42)$$

4. Global energy and angular momentum

If we now consider the particular case of a stationary and/or axisymmetric system, in which there will be a conserved energy and/or angular momentum per particle given by

$$\mathcal{E}^x = -\vec{k} \cdot \underline{\pi}^x \quad (43)$$

$$\mathcal{J}^x = \vec{m} \cdot \underline{\pi}^x \quad (44)$$

then we shall have

$$\mathcal{P}_{\text{MC}}^a \{k\} d\Sigma_a = -\mathcal{E}^x dN_x = \sum_x \vec{\mathcal{E}}_{\text{CM}}^x \cdot d\underline{\Sigma} \quad (45)$$

$$\mathcal{P}_{\text{MC}}^a \{m\} d\Sigma_a = \mathcal{J}^x dN_x = -\sum_x \vec{\mathcal{J}}_{\text{CM}}^x \cdot d\underline{\Sigma} \quad (46)$$

across any respectively stationary or axisymmetric hypersurface, and in particular across a black hole horizon, where we define the fluxes

$$\vec{\mathcal{E}}_{\text{MC}}^x = \mathcal{E}^x \vec{n}_{|x|} \quad (47)$$

$$\vec{\mathcal{J}}_{\text{MC}}^x = \mathcal{J}^x \vec{n}_{|x|} \quad (48)$$

which evidently satisfy the separate conservation laws

$$\nabla \cdot \vec{\mathcal{E}}_{\text{MC}}^x = 0 \quad (49)$$

$$\nabla \cdot \vec{\mathcal{J}}_{\text{MC}}^x = 0 . \quad (50)$$

In the case of angular momentum, the total exterior source contribution given by (6.322)¹ may be expressed by

$$J_{\text{MC}} = \sum_x \int_{\Sigma} \vec{\mathcal{J}}_{\text{MC}}^x \cdot d\underline{\Sigma} \quad (51)$$

for any axisymmetric hypersurface Σ , but the fact that Σ could not be invariant under $\vec{k}$ prevents us from obtaining an equally simple formula for M_{MC} .

We now proceed from (6) as in Section 6.6.1 using the strict stationary boundary conditions (6.314)¹ to evaluate the surface integrals on the horizon in the form

$$\delta \oint_S \frac{1}{2} k^c A_c D_{\text{F}}^{ba} dS - \oint_S D_{\text{F}}^{c[b} (\delta A_c) k^{a]} dS_{ab} = \Phi^{\text{H}} \delta Q_{\text{H}} + \Omega^{\text{H}} \delta J_{\text{FH}} \quad (52)$$

We next perform the analogous steps in the evaluation of the material contributions, using the formulae of the previous section which enable us to write firstly

$$-II_{\text{M}}^a\{k\} = (\Lambda_{\text{M}} - n_x^b \mu_b^x) k^a + (k^b \mu_b^x) n_x^a \quad (53)$$

and

$$|g|^{-1/2} \delta(|g|^{1/2} \Lambda_{\text{M}}) = \frac{1}{2} (T_{\text{M}c}^b h_b^c + n_x^b \mu_b^x h_c^c) + \mu_b^x \delta n_x^b. \quad (54)$$

Since we now only have source terms to deal with we proceed directly (i.e. without recourse to Green's theorem) using (6.399)¹ to obtain

$$-\delta \int_{\Sigma} II_{\text{M}}^a d\Sigma_a - \frac{1}{2} \int_{\Sigma} T_{\text{M}c}^b h_b^c k^a d\Sigma_a = \int_{\Sigma} k^b \mu_b^x \delta(n_x^a d\Sigma_a) + \frac{1}{2} \int_{\Sigma} k^{[b} n_x^{a]} (\delta \mu_b^x) d\Sigma_a \quad (55)$$

in which the terms can be seen to be closely analogous to the volume integral terms in (6). Thus using the expression (12) for the *total momenta* we can combine (6) and (55) so as to obtain (with the aid of (52) and (6.320)¹) the final result

$$\delta M - \Omega^{\text{H}} \delta J_{\text{BH}} - \Phi^{\text{H}} \delta Q_{\text{H}} - \frac{\kappa}{8\pi} \delta \mathcal{A} = \int_{\Sigma} \vec{k} \cdot \underline{\pi}^x \delta(\vec{n}_x \cdot d\underline{\Sigma}) + \int_{\Sigma} (\delta \underline{\pi}^x) \cdot (\vec{k} \wedge \vec{n}_x) \cdot d\underline{\Sigma} \quad (56)$$

(in which $J_{\text{BH}} = J_{\text{H}} + J_{\text{FH}} = J - J_{\text{MC}}$).

5. Interpretation in the generic (convective) case

The first term on the right hand side of (56) can be interpreted as representing the *static* injection energy: it has the form

$$\int_{\Sigma} \vec{k} \cdot \underline{\pi}^x \delta(\vec{n}_x \cdot d\underline{\Sigma}) = \int \mathcal{E}^x \delta(dN_x) \quad (57)$$

where dN_x , as given by (42), is the flux of the x th constituent across $d\underline{\Sigma}$, and where $\mathcal{E}^x$ is the canonical *energy per particle*, as given by (43), which satisfies the conservation law

$$\vec{n}_{|x|} \cdot \nabla \mathcal{E}^x = 0 \quad (58)$$

for each x .

The form of the second term in (56) (which vanishes altogether in a static situation, i.e. one with all current vectors parallel to $\vec{k}$) may seem less familiar, but it is in fact a generalisation of the Faraday term $I\delta\mathcal{F}$ representing the energy needed to make a change $\delta\mathcal{F}$ in the magnetic flux $\mathcal{F}$ through a circuit carrying a current I . The relationship can be made rather more apparent if we decompose each flux $\vec{n}_x$ into a *static* part plus a spacelike part $\vec{c}_x$ that is confined within the spacelike hypersurface Σ , in the form

$$\vec{n}_x = n_x^{(0)} \vec{k} + \vec{c}_x, \quad \vec{c}_x \cdot d\underline{\Sigma} = 0 \quad (59)$$

since we may then write

$$\int_{\Sigma} (\delta \underline{\pi}^x) \cdot (\vec{k} \wedge \vec{n}_x) \cdot d\underline{\Sigma} = - \int_{\Sigma} (\vec{c}_x \delta \underline{\pi}^x) (\vec{k} \cdot d\underline{\Sigma}). \quad (60)$$

We are always at liberty to decompose the hypersurface element $d\underline{\Sigma}$ in the form

$$d\underline{\Sigma} = d\vec{x} \cdot d\underline{\mathcal{S}} \quad (61)$$

where $d\vec{x}$ is an arbitrary infinitesimal displacement and $d\underline{\mathcal{S}}$ is a normal 2 surface element transverse to the displacement. If we take $d\vec{x}$ to be locally parallel to one of the fluxes, $\vec{c}_x$ say, we shall have

$$-(\vec{c}_{|x|} \delta \pi^x) (\vec{k} \cdot d\underline{\Sigma}) = (d\vec{x} \cdot \delta \pi^x) dC_{|x|} \quad (62)$$

where $dC_{|x|}$ represents the current of the x th constituent through the element $d\underline{\mathcal{S}}$ as defined by

$$dC_x = \vec{c}_x \cdot (\vec{k} \cdot d\underline{\mathcal{S}}) . \quad (63)$$

Thus dC_x represents the number of particles crossing the element $d\underline{\mathcal{S}}$ per unit of the *ignorable* (not proper) group time t . In general one can envisage that the flux lines in Σ will have complicated ergodic properties. However *if* their global structure is such that they form sufficiently well behaved *simple closed circuits* then one can see from (62) that we shall have

$$\delta M - \Omega^H \delta J_{\text{BH}} - \Phi^H \delta Q_H - \frac{\kappa}{8\pi} \delta \mathcal{A} = \int \mathcal{E}^x \delta(dN_x) + \int (\delta \mathcal{C}^x) dC_x \quad (64)$$

where for each circuit the conserved circulation integral $\mathcal{C}^x$, which is a generalisation of the Faraday flux, is as defined by (24).

6. The circular (non - convective) case

Despite the restriction imposed by the non-ergodicity hypothesis, the deceptively simple form (64) still represents quite a powerful generalisation compared not only with the original variation law of Bardeen *et al* (1973)^{2,3}, but even with the electromagnetic extension of Carter (1973b)⁴, since these earlier versions applied only to flows with strictly (not just topologically) circular circuits. To show how these earlier versions are obtained in the appropriate, non - convective (and non - conducting) limit, we first introduce a vector $\vec{v}_x$ defined by

$$\vec{c}_x = n_x^{(0)} \vec{v}_{|x|} \quad (65)$$

and representing the mean *displacement* of the particles of the x th constituent per unit of the *ignorable* (not proper) group time t . (If there were any particles in the immediate neighbourhood of the black hole, their 3-velocities as thus defined would evidently have to tend towards the Damour⁷ velocity $\vec{v}_H$ – as defined in section 6.3¹ – in the limit on the horizon.) We see that in terms of the corresponding unit flow vectors $\vec{u}_x$, defined by

$$\vec{n}_x = n_x \vec{u}_{|x|} , \quad |\vec{u}_{|x|}|^2 = -1 \quad (66)$$

the 3-velocity 4-vectors will be given by

$$\vec{k} + \vec{v}_x = \dot{\tau}_{|x|} \vec{u}_x \quad (67)$$

where $\dot{\tau}_x$ is the relevant time dilation (or “mean redshift”) factor. We can use these 3-velocities to regroup the terms on the right hand side of (56) so as to obtain the variation law

$$\delta M - \Omega^H \delta J_{\text{BH}} - \Phi^H \delta Q_H - \frac{\kappa}{8\pi} \delta \mathcal{A} = \int_{\Sigma} \pi_{(0)}^x \delta(dN_x) - \int_{\Sigma} \vec{v}_x^b \delta(\pi_{,b}^x n_x^a d\Sigma_a) \quad (68)$$

where

$$\pi_{(0)}^x = -(\vec{k} + \vec{v}_{|x|}) \cdot \underline{\pi}^x \quad (69)$$

which is valid quite generally (regardless of the topological behaviour of the flow lines).

If we now impose the restriction that the flow lines be strictly *circular*, so that the 3-velocities take the form

$$\vec{v}_x = \Omega_x \vec{m} \quad (70)$$

where Ω_x are the corresponding *angular velocities* then we shall have

$$\delta M - \Omega^H \delta J_{\text{BH}} - \Phi^H \delta Q_H - \frac{\kappa}{8\pi} \delta \mathcal{A} = \int_{\Sigma} \pi_{(0)}^x \delta(dN_x) + \int_{\Sigma} \Omega_x \delta(dJ_{\text{MC}}^x) \quad (71)$$

with

$$dJ_{\text{MC}}^x = \vec{\mathcal{J}}_{\text{MC}}^x \cdot d\Sigma \quad (72)$$

where the constituent angular momentum fluxes are given by (48). The quantity $\pi_{(0)}^x$, which is a generalisation of what Bardeen² has called the zero angular momentum injection energy, is given by

$$\pi_{(0)}^x = \mathcal{E}^x - \Omega_{|x|} \mathcal{J}^x = \mu_{(0)}^x + e^x \Phi_{|x|(0)} \quad (73)$$

where the *effective* chemical potential $\mu_{(0)}^x$ is related to the corresponding local rest frame chemical potential μ^x by

$$\mu_{(0)}^x = -(\vec{k} + \vec{v}_{|x|}) \cdot \underline{\mu}^x = \dot{\tau}_{|x|} \mu^x \quad (74)$$

with

$$\mu^x = -\vec{u}_{|x|} \cdot \underline{\mu}^x \quad (75)$$

and where the corotating electric potential $\Phi_{x(0)}$ is given by

$$\Phi_{x(0)} = -(\vec{k} + \vec{v}_{|x|}) \cdot \underline{A} = \dot{\tau}_{|x|} \Phi_x \quad (76)$$

with

$$\Phi_x = -\vec{u}_{|x|} \cdot \underline{A} . \quad (77)$$

References.

1. B. Carter, “The general theory of the mechanical, electromagnetic and thermodynamical properties of black holes”, in *General Relativity: an Einstein centenary survey*, ed S.W. Hawking & W. Israel *pp* 295 - 369 (Cambridge U.P. 1979).
2. J.M. Bardeen, “Rapidly rotating stars, discs and black holes”, in *Black Holes*, ed B. DeWitt & C. DeWitt *pp* 215 - 289 (Gordon & Breach, New York, 1973).
3. J. Bardeen, B. Carter, & S.W. Hawking, “The Four Laws of Black Hole Mechanics”, *Commun. Math. Phys.* **31**, *pp* 161-170 (1973).
4. B. Carter, “Black Hole Equilibrium States: II General Theory of Stationary Black Hole States”, in *Black Holes* (proc. 1972 Les Houches Summer School), ed. B. De Witt & C. DeWitt, *pp* 125-210 (Gordon & Breach, New York, 1973).
5. B. Carter, “Regular and Anomalous Heat Conduction: the Canonical Diffusion Equation in Relativistic Thermodynamics”, in *Journées Relativistes 1976*, ed. M. Cahen, R. Debever, & J. Geheniau, *pp* 12-27 (Université Libre de Bruxelles, 1976).
6. B. Carter, “Elastic Perturbation Theory in General Relativity and a Variational Principle for a Rotating Solid Star”, *Commun. Math. Phys.* **30**, *pp* 261-286 (1973).
7. T. Damour, “Black hole eddy currents”, *Phys. Rev.* **D18**, *pp* 3598-3604 (1978).